


## Effect of Light and Heavy Smoking on High Density Lipoprotein Cholesterol Level

Haydar Al-Shalah

College of Medicine, University of Babylon, Hilla, Iraq.



### Abstract

**Background and purpose:** Cigarette smokers have a higher risk of developing several chronic disorders. These include several types of cancer, chronic obstructive pulmonary disease and atherosclerosis which is a chief contributor to the high number of deaths from smoking. Cigarette smoking potentiates the harmful effects of total cholesterol and reduces the cardio-protective properties of high density lipoprotein cholesterol (HDLc), increasing the risk of coronary heart disease(CHD). Decreased serum HDLc is one of the most common lipid disorders in patients with CHD. Existing evidence suggests that every 1 mg/dl decrease in serum HDLc increases the risk of CHD by 2-3%.

HDLc levels are known to be lower in smokers than in nonsmokers. However most studies have not been mentioned whether heavy or light smoking affect HDLc level. The aim of the present study was to evaluate the effect of light and heavy smoking on HDLc level.

**Material and method:** Sixty apparently healthy participants were included in the study, twenty subjects as control group, twenty light smokers(those who smoke less than ten cigarettes per a day) and twenty heavy smokers (those who smoke ten cigarettes and more per a day).Their mean age  $\pm$  SD was  $35.25\pm 4.38, 37.25\pm 4.74, 37\pm 3.93$  for control, light smokers and heavy smokers groups respectively .The mean of the BMI of the same groups was  $23.36\pm 1.7, 22.91\pm 1.67$  and  $23\pm 1.49$ .HDL-C was measured by colorimetric method.

**Results:** The mean values of HDLc of control ,light smokers and heavy smokers was  $1.43\pm 0.42$  mmol/l,  $0.75\pm 0.17$  mmol/l. and  $0.6\pm 0.2$  mmol/l groups respectively .There was significant differences in the mean level of HDLc between control group and light smoker and heavy smoker groups( $P < 0.05$ ).However the mean value of HDLc of light smokers was not significantly different from the mean level of heavy smokers( $P > 0.05$ ).

**Conclusion:** Smoking whether light or heavy equally affect HDLc level by decreasing its concentration in the serum and hence increase the risk of CHD.

### تأثير التدخين الخفيف والتدخين المفرط على مستوى الكوليسترول الحميد

#### الخلاصة

المدخنين هم الأكثر عرضة للإصابة بالسرطانات على مختلف أنواعها وأمراض الرئة بالإضافة إلى تصلب الشرايين والأخير هو السبب الرئيسي للموت من التدخين. التدخين يؤدي إلى زيادة نسبة الكوليسترول وتقليل نسبة الكوليسترول الحميد وبالتالي يؤدي إلى أمراض القلب التاجية كون الأخير له صفة الحماية للقلب من أمراض الشرايين.

ان معظم الدراسات تشير إلى أن التدخين يؤدي إلى نقصان مستوى الكوليسترول الحميد في حين لم تشير تلك الدراسات إلى أي نوع من التدخين (الخفيف والمفرط) يؤدي إلى ذلك النقصان .

أجريت هذه الدراسة في مدينة بابل في أيلول من العام 2009 لغرض تقييم مستوى الكوليسترول الحميد في أمصال 20 من المدخنين الذين يدخنون اقل من عشرة سكاثر في اليوم الواحد و20 من المدخنين الذين يدخنون عشرة سكاثر فما فوق بالإضافة إلى 20 شخصا من غير المدخنين كمجموعة مسيطرة .

كان معدل أعمار المجموعات الثلاث 35.25 و37.25 و37 سنة على التوالي. كان معدل الكوليسترول الحميد للمدخنين المفرطين 0.6 مل مول للتر و المدخنين الذين يدخنون أقل من 10 سكاثر في اليوم 0.75 مل مول للتر و 1.43 مل مول للتر لغير المدخنين. كان هناك علاقة ذات دلالة عالية (مستوى الاحتمالية أقل من 0.05) بين معدل مستوى الكوليسترول الحميد بين غير المدخنين والمدخنين المفرطين والمدخنين الخفيف في حين لم تكن هناك علاقة ذات دلالة عالية (مستوى الاحتمالية أكثر من 0.05) بين المدخنين من كلا المجموعتين وهذا يدل على أن التدخين الخفيف والمفرط يؤدي إلى انخفاض نسبة الكوليسترول الحميد وبشكل متساو.

## Introduction

Cigarette smoking are social habits of lifestyle that are closely related to a variety of diseases. Smoking is a major risk factor for atherosclerotic cerebro- and cardiovascular diseases. Various pathophysiological mechanisms, including injury of the vascular endothelium and lipid peroxidation, have been known to be involved in facilitation of the atherosclerotic progression due to smoking [1,2]. Smoking affects the blood lipid profile: serum HDL and LDL cholesterol concentrations have been reported to be lower and higher, respectively, in smokers than in non-smokers [3,4]. Thus, smoking may exert its atherogenic effects through blood lipid levels [5,6].

High density lipoprotein (HDL) is a plasma lipoprotein transporting 30.0% phospholipids, 25.0% cholesterol and 5.0% triacylglycerol. Its high density is due to 40.0% proteins. The cholesterol associated with it is unutilized excess cholesterol from tissues. This cholesterol is returned to liver. The liver converts it into bile acids, bile salts, and cholesteryl esters. The two bile salts formed in the liver are sodium glycocholate and sodium taurocholate. These compounds are secreted by liver into bile and through bile they reach the small intestines. The returning of the scavenged cholesterol to liver from peripheral tissues by HDL is known as "Reverse cholesterol transport"[7].

Pre- $\beta$ -HDL cholesterol; rich in apoprotein A-1 is synthesized by liver and intestinal mucosal cells and released into circulation. Plasma lecithin-

cholesterol acyl transferase (LCAT) converts free cholesterol in pre- $\beta$ -HDL to cholesteryl ester, resulting in the formation of  $\alpha$ -HDL cholesterol. LCAT catalyzes the freely reversible reaction which transfers the fatty acid at C2 in lecithin to cholesterol. It is  $\alpha$ -HDL which takes part in the reverse cholesterol transport. HDLc inhibits the oxidation of low density lipoprotein cholesterol (LDLc) and reduce the risk of thrombosis by inhibiting platelet aggregation[8].

A higher level of HDLc is an index of safety from the risk of coronary heart Disease (CHD). This is due to the fact that unutilized excess cholesterol which is potentially harmful, is scavenged from the tissues by HDL preventing its deposition and plaque formation[9]. Low plasma HDLc is a risk factor for CHD particularly in the male[10,11]. Experimental studies in laboratory animals and epidemiological studies in population groups have shown that raising the level of HDLc in plasma may retard the development of atherosclerosis[12].

Effect of life style modification on HDLc in plasma are many such as Exercise; walking seems to have no significant effect in increasing HDLc in plasma[13]. Regular aerobic exercises increase HDLc in plasma up to 9.0% in sedentary people[14]. Exercise is believed to increase HDLc by stimulating the production of pre- $\beta$ -HDLc and increasing reverse cholesterol transport[15]. Cigarette smoking is associated with lowered HDLc and LCAT activity[16,17]. After cessation of smoking HDLc is observed to

increase[18].Moderate alcohol intake is good as it raises the plasma HDLc by increasing esterified fraction of cholesterol in plasma[19]. Both LDLc and HDLc in plasma are reduced with low intake of fat. A fat rich in polyunsaturated fatty acids has a beneficial effect in raising plasma HDLc[20].

Medication such as Niacin, Fibrates and Statins have been found to increase HDLc in plasma. These medications also lower LDLc and TG in plasma. Niacin is most effective in raising plasma HDLc. However, the side effects of niacin like cutaneous flushing and dyspepsia have to be taken care of. Fibrate therapy has resulted in an increase of plasma HDLc by 10-25%[21].Gemfibrozil has significantly reduced the risk of coronary heart disease. Statins (Simvastatin,Atorvastatin) have been found to elevate plasma HDLc by raising apolipoprotein A-1 synthesis[22].

**Aim of the Study**

HDLc levels are known to be lower in smokers than in nonsmokers. However most studies have not been mentioned whether heavy or light smoking affect HDLc level. The aim of the present study was to evaluate the effect of light and heavy smoking on HDLc level.

**Material and Methods**

The study was conducted in Babylon city during September 2009. The study included 60 apparently healthy male subjects who are similar in terms of age and physical activity. Twenty heavy smokers(those who smoke 10 cigarettes and more per a day). Twenty light smokers( those who smoke less than 10

cigarettes per a day ). Another 20 subject who are non smokers included in this study as a control group. The duration of smoking of the first two group was between five and ten years.

All subjects were required to meet certain criteria. None of the subjects drank alcohol or taking medication. None of them were obese (<25kg/m<sup>2</sup>).All subjects fasted for 14-16 hours before their blood was drawn to measure the serum HDL. Body weight and height were measured and the BMI was calculated as weight/height<sup>2</sup> (kg/m<sup>2</sup>).

HDL-C concentrations were measured by enzymatic techniques in the biochemistry laboratory of the medical college. HDLc levels were measured after precipitation with phosphotungstic acid and magnesium. Kits for the HDLc assays were from Biolabo (France).

Statistical analysis: SPSS (version 10) program was used for statistical analysis. The results were represented through mean and standard deviation. F-test was used to compare between means & to study the significance of the difference. P value < 0.05 was considered to be statistically significant.

**Results**

A total of sixty apparently healthy male subjects were studied, twenty heavy smokers,twenty light smokers in addition to twenty non smokers were enrolled for comparative assessment. Their mean age ±SD was 37±3.93 ,37.25±4.74,35.25±4.38.(table 1).The BMI of the same groups was23±1.49 ,22.91±1.67 and 23.36±1.7 respectively.(table1)

**Table 1** Numbers ,age and body mass index in heavy smokers, light smokers and non smokers groups

| Subject characteristics       | Heavy smokers | Light smokers | Non smokers |
|-------------------------------|---------------|---------------|-------------|
| Number                        | 20            | 20            | 20          |
| Age(mean±SD)years             | 37±3.93       | 37.25±4.74    | 35.25±4.38  |
| BMI(mean±SD)kg/m <sup>2</sup> | 23±1.49       | 22.91±1.67    | 23.36±1.7   |

There was no significant differences in terms of age and BMI between all three groups(P value >0.05).

The mean value of HDLc of non smokers was 1.43±0.42 mmol/l which was significantly different(P value <0.05) from mean value of HDLc of

heavy smokers(0.6±0.2 mmol/l) and light smokers group(0.75±0.17 mmol/l.).However ,there was no significant differences in the mean value of HDLc between light and heavy smoker groups(P value >0.05).(table2)

**Table 2** Mean value of serum level of HDLc of non smokers, light smokers and heavy smokers groups

| Group         | Mean value of HDLc ±SD(mmol/l) |
|---------------|--------------------------------|
| Non smokers   | 1.43±0.42                      |
| Light smokers | 0.75±0.17                      |
| Heavy smokers | 0.6±0.2                        |

**Discussion**

It is well known that HDLc is inversely correlated with the risk of atherosclerosis and coronary heart disease .Similarly, it is well known that smoking affect HDLc by deceasing its level in the serum, therefore this study was carried out to shade a light on which type of smoking heavy or light decrease HDLc level.

The protective effect of HDL has been related to their role in the cholesterol reverse transport (from peripheral tissues to the liver) and to their ability to inhibit oxidation of low-density lipoproteins (LDL) and biological membranes [8,9]. Smoking affects the blood lipid profile: serum HDL and LDL cholesterol concentrations have been reported to be lower and higher, respectively, in smokers than in non-smokers .Thus, smoking may exert its atherogenic effects through blood lipid levels [3-6].

It has been reported that the safest level of HDLc to be protective against CHD is more than 1.17 mmol/l[7,9,10,11]. In the present study, the mean value of HDLc of control group was 1.43mmol/l.

Furthermore,in the present study HDLc was low in smokers compared with control group and it is identical with most of the studies[4,17,18]. However its level was low in both

groups of smokers whether light or heavy.

**Conclusion**

Smoking whether light or heavy equally affect HDLc level by deceasing its concentration in the serum and hence increase the risk of CHD.

**References**

- 1-Tsiara S, Elisaf M , Mikhailidis D P. Influence of smoking on predictors of vascular disease. *Angiology* 2003 ; 54: 507–30.
- 2- Ambrose J A, Barua RS .The pathophysiology of cigarette smoking and cardiovascular disease: An update. *Journal of the American College of Cardiology* 2004 ;43: 1731–37.
- 3- Whitehead T P, Robinson D, Allaway S L. The effects of cigarette smoking and alcohol consumption on blood lipids: Adose-related study on men. *Annals of Clinical Biochemistry* 1996; 33: 99–106.
- 4- Lee K S, Park CY, Meng K H;et al. The association of cigarette smoking and alcohol consumption with other cardiovascular. *Lancet*2000 ; 96:1965-8.
- 5- Corrao G , Rubbiati L, Bagnardi V, Zambon A, Poikolainen K. Alcohol and coronary heart disease: A meta-analysis. *Addiction* 2000;95: 1505–23.
- 6- Meister K A, Whelan E M, Kava R. The health effects of moderate alcohol

intake in humans: An epidemiologic review. *Critical Reviews in Clinical Laboratory Sciences* 2000; 37: 261–96.

7. Brewer HB. High Density Lipoproteins: a new potential therapeutic target for the prevention of cardiovascular disease. *Arterioscler Thromb Vasc Biol* 2004; 24: 387-91.

8- Shah PK, Kaul S, Nilsson J, Cercek B. Exploiting the vascular protective effects of high density lipoprotein and its apolipoproteins: an idea whose time for testing is coming. *Circulation* 2001; 104: 23783.

9. Rajagopal G. Cholesterol-an update. *Biomed* 2003; 23: 1-8.

10. Miller NE, Thelle DS, Forde OH, Mijos OD. High density lipoprotein and coronary heart disease: a prospective case control study. *Lancet* 1997; 66:965-8.

11. Wilson PW, Abbott RD, Castelli WP. High density lipoprotein cholesterol and mortality: The Framingham heart study. *Arteriosclerosis* 1988; 8: 737-41.

12-Gordon DJ, Knoke J, Probstfield JL, Superko R, Tyroler HA. High density lipoprotein cholesterol and coronary heart disease in hypercholesterolemic men :the lipid research clinics coronary primary prevention trial. *Circulation* 1986; 74: 1217-25.

13- Kelly GA, Kelly KS, Tran ZV. Walking, lipids and lipoproteins: a meta analysis of randomized controlled trials. *Preventive Med* 2004; 38: 651-61.

14- King AC, Haskell WI, Young DR, Oka RK, Stefanick ML. Long term effects of varying intensities and formats of physical activity on participation rates, fitness, and lipoproteins in men and women aged 50-65 years. *Circulation* 1995; 91: 2596-604.

15- Gupta AK, Ross EA, Myels JN, Kashyap L. Increased reverse cholesterol transport in athletes. *Metabolism* 1993; 42: 684-90.

16- Ellison RC, Zgang Y, Qureshi MH, Knox S, Arnett DK. Life style

determinants of high density lipoprotein cholesterol: the National Heart, Lung and Blood Institute Family Heart Study. *Amer Heart J* 2004; 147: 529-35.

17- Inamura H, Teshima K, Miyamoto N, Shirota T. Cigarette smoking, high density lipoprotein cholesterol sub-fractions and lecithin:cholesterol acyl transferase in young women. *Metabolism* 2002; 51: 1313-6.

18- Marda K, Noguchi Y, Fukui T. The effects from cessation from cigarette smoking on the lipid and lipoprotein profiles: a meta analysis. *Preventive Med* 2003; 37: 283-90.

19. Van der Gagg MS, Vantel A, Vermunt SH, Scheek LM, Schaafsma G, Hendricks HF. Alcohol consumption stimulates early steps in reverse transport of cholesterol. *J Lipid Res* 2001; 42: 2077-83.

20. Berglund L, Oliver EH, Fontanez N ;*et al.* HDL sub-population patterns in response to deductions in dietary total unsaturated fat intakes in healthy subjects. *Amer J Clin Nutr* 1999; 70: 992-1000.

21. Dominique M, Roger S. Low HDL cholesterol levels. *New Engl J Med* 2005; 353: 1252-60.

22. Schaefer JR, Schweer H, Ikewaki K; *et al.* Metabolic basis of high density lipoproteins and apolipoproteins A-1 increase by HMG CoA reductase inhibition in healthy subjects. *Atherosclerosis* 1999; 144: 177-84.