

On Completely And Semi Completely Prime Ideal With Respect To An Element Of A Boolean α_1, α_2 Near-Ring

Ameer A. J. AL Swidi

Abstract: In this paper ,I generalize concepts of α_1, α_2 near-ring as well as of completely semi prime ideal with respect to an element x of a α_1, α_2 near-ring and the completely prime ideal of a α_1, α_2 near-ring with respect to an element x and the relationships between the completely prime ideal with respect to an element x of a α_1, α_2 near-ring and some other types of ideals,as well as I will valid if I is c.p.I of α_1, α_2 near-ring iff is a x-c.s.p.I of a α_1, α_2 near-ring as well as valid if I is c.p.I of α_1, α_2 near-ring iff is a x-c.s.p.I of α_1, α_2 near-ring as well as I is e'-c.p.I of α_1, α_2 near-ring iff it is a e'-c.s.p.I of α_1, α_2 near-ring and all its valid after put condition of Boolean a α_1, α_2 near-ring

Keywords : α_1, α_2 near-ring, c.s.p.I, c.p.I, x-c.s.p.I, x-c.p.I

1 INTRODUCTION

In 1905 L.E Dickson began the study of near- ring and later 1930 Wieland has investigated it .Further, material about a near ring can be found (cf.[2,5,9]).In 1977 G. Pilz, was introducing the notion of prime ideals of a near-ring (cf.[1,2,7]).. In 1988 N.J.Groenewald was introducing the notions of completely (semi) prime ideals of a near-ring (cf.[6,10]). In 2011 H.H.abbass, S.M.Ibrahem introduced the concepts of a completely semi prime ideal with respect to an element of a near-ring[4] .In 2012 H,H.abbass,Mohanad Ali Mohammed introduce the concept of a completely semi prime ideals near-ring with respect to an element of a near-ring [3]. In2010,S.Uma,R.Balakrishnan,T.Tamizh Chelvam introduce the concept of a α_1, α_2 near-ring near- ring[8].

1.1 definition

A right near-ring is a set N together with two binary operations + and . such that

1. (N,+) is a group (not necessarily abelian)
2. (N, .) is a semigroup
3. $n_1.(n_2+n_3) = n_1 . n_2 + n_1 . n_3 , \forall n_1, n_2, n_3 \in N$

1.2 definition:

Let N be a right near-ring , if

1-for every a in N there exists x in N such that $a=xax$ then we say N is an α_1 near-ring.

2- for every a in $N-\{0\}$ there exists x in $N-\{0\}$ such that $xax=x$ then we say N is an α_2 near-ring.

1.3 example:

1-the near-ring (N,+.) define on the klein's four group $N=\{0,a,b,c\}$ where addition and multiplication is defined as:

+	0	a	b	c		.	0	a	b	c
0	0	a	b	c		0	0	0	0	0
a	a	0	c	b		a	0	0	a	a
b	b	c	0	a		b	0	a	b	b
c	c	b	a	0		c	0	a	c	c

Is an α_1 near-ring (since $bab=ab=a$, $bbb=b$, $ccc=c$, $a0a=0$) its neither α_2 near-ring (since there is no x in $N-\{0\}$ such that $xax=x$) 2-the near-ring (N,+.) define on the klein's four group $N=\{0,a,b,c\}$,where addition and multiplication is defined as:

+	0	a	b	c		.	0	a	b	c
0	0	a	b	c		0	0	0	0	0
a	a	0	c	b		a	a	a	a	a
b	b	c	0	a		b	0	0	b	0
c	c	b	a	0		c	a	a	c	a

Here (N,+.) is an α_2 near-ring it is neither α_1 near-ring (since $xcx \neq c$ for any x in N).

1.4 definition:

Let N be a α_1, α_2 near-ring , normal subgroup I of (N,+) is called a right ideal of N if :

1- $NI \subseteq I$,Where $NI=\{n.i : n \in N , i \in I\}$

2- $\forall n, n_1 \in N$ and for all $i \in I$, $(n_1+i)n-n_1n \in I$

1.5 example:

Consider the a N be a α_1, α_2 near-ring example(1.3) the normal subgroup $I=\{0,a\}$ is ideal of the near-ring N .

1.6 Remark

We will refer that all ideals in this paper are right ideals.

- Ameer A. J. AL Swidi
- University of Babylon, College of Education for pure science. Mathematical Department

1.7 definition:

Let N be a α_1, α_2 near-ring, an ideal I of N is called a completely semi prime ideal (c.s.p.I) of N if $x^2 \in I$ implies $x \in I$ for all $x \in N$.

1.8 example:

Let N be a α_1, α_2 near-ring, the ideal $I = \{0, a\}$ of N in example(1.3) is a completely semi prime ideal of N, since $\forall y \in N$ implies $y \in I$.

1.9 definition:

Let N be a α_1, α_2 near-ring, an ideal of N is called a prime ideal if every ideal I_1, I_2 of N such that $I_1 \cdot I_2 \subseteq I$ implies $I_1 \subseteq I$ or $I_2 \subseteq I$.

1.10 example:

Let N be a α_1, α_2 near-ring, in example(1.3) the ideal $I = \{0, a\}$ of the N is a prime ideal.

1.11 definition:

Let N be a α_1, α_2 near-ring, s be an ideal in N so s is semi prime if and only if for all ideals I of N, $I^2 \subseteq s$ implies $I \subseteq s$.

1.12 definition:

Let N be a α_1, α_2 near-ring, I be an ideal of N, then I is called a completely prime ideal of N if for all $x, y \in N$, $x \cdot y \in I$ implies $x \in I$ or $y \in I$ denoted by (c.p.I) of N.

1.13 example

Let N be a α_1, α_2 near-ring, in example(1.3) the ideal $I = \{0, a\}$ is complete prime ideal of the N.

1.14 proposition:

Let N be a α_1, α_2 near-ring, every c.p.I of N is a c.s.p.I of N.

Proof:

Let for all $y \in N$, such that

$$y^2 \in I \text{ as } I \text{ is c.p.I of } N$$

$$\text{So, } y^2 = y \cdot y \in I$$

then, $y \in I$

so, I is a c.s.p.I of N.

1.15 definition:

Let N be a α_1, α_2 near-ring, is called Boolean if for all $x \in N, x^2 = x$.

1.16 example:

the near ring $(N, +, \cdot)$ define on the Klein's four group $N = \{0, a, b, c\}$, where addition and multiplication is defined as :

+	0	a	b	c	.	0	a	b	C
0	0	a	b	c	0	0	0	0	0
a	a	0	c	b	a	a	a	a	a
b	b	c	0	a	b	0	0	b	b
c	c	b	a	0	c	a	a	c	c

Is a α_1, α_2 near-ring, as every Boolean near ring is α_1 near-ring as well as α_2 near-ring since $(00=00=0, aaa=aa=a, bbb=bb=b, ccc=cc=c)$.

2 Completely and semi completely prime ideal with respect to an element of a α_1, α_2 near-ring

2.1 definition:

Let N be a α_1, α_2 near-ring and $x \in N$, I is called completely semi prime ideal with respect to an element x denoted by (x-c.s.p.I) of N if for all $y \in N$, $x \cdot y^2 \in I$ implies $y \in I$.

2.2 example:

Consider the near ring $N = \{0, a, b, c\}$, where addition and multiplication is defined as :

+	0	a	b	c	.	0	a	b	c
0	0	a	b	c	0	0	0	0	0
a	a	0	c	b	a	0	a	0	a
b	b	c	0	a	b	0	0	b	b
c	c	b	a	0	c	0	a	b	c

is a α_1, α_2 near-ring, the ideal $I = \{0, b\}$ is a-c.s.p.I of N.

2.3 definition:

Let N be a α_1, α_2 near-ring, I be an ideal of N and $x \in N$, then is called a completely prime ideal with respect to an element x denoted by (x-c.p.I) of N if for all $y, z \in N$, $x \cdot y \cdot z \in I$ implies $y \in I$ or $z \in I$.

2.4 example:

Let N be a α_1, α_2 near-ring, in example(2.2) the ideal $I = \{0, b\}$ is a-c.p.I of N.

2.5 proposition:

Let N be a α_1, α_2 near-ring and $x \in N$, then every completely prime ideal with respect to an element x of N is a completely semi prime ideal with respect to an element x of N.

Proof:

let I be a x-c.p.I and if for all $y \in N$ such that $x \cdot y^2 \in I$,

$$x \cdot y^2 = x \cdot y \cdot y \in I$$

implies $y \in I$

then, I is x-c.s.p.I of N.

2.6 Remark:

The converse of the proposition (2.5) may be not true, just where put the condition of Boolean of N its valid (proposition(2.11)).

2.7 proposition:

Let N be a Boolean α_1, α_2 near-ring then I be c.p.I of N iff I is an x-c.s.p.I for all $x \in N - I$.

Proof:

(\Rightarrow) Let $x, y \in N$ such that $x.y^2 \in I$ as N is Boolean, then $x.y^2 \in I \rightarrow x.y \in I$ so $y \in I$ as I is c.p.I of N then I is an x-c.s.p.I for all $x \in N-I$
 (\Leftarrow) let $\forall y \in N, x.y \in I$ then $x.y^2 \in I$ so that $y \in I$ then I is c.p.I of N .

2.8 proposition:

Let N be a Boolean α_1, α_2 near-ring, then I be c.p.I of N iff I is an x-c.p.I for all $x \in N$.

Proof:

(\Rightarrow) as I is c.p.I of α_1, α_2 near-ring Let $\forall y \in N, x.y.y \in I$ then $\forall y \in N, x.y^2 \in I$ as N is Boolean, then $\forall y \in N, x.y \in I$ and so $y \in I$ as I is c.p.I of N then I is x-c.p.I of N .
 (\Leftarrow) let $\forall y \in N, x.y \in I$ as N is Boolean, then $x.y^2 \in I \rightarrow x.y.y \in I$ so $y \in I$, as I is an x-c.p.I of N then, I is c.p.I of N .

2.9 Proposition:

Let N be a Boolean α_1, α_2 near-ring, then every x-c.p.I of N is c.s.p.I of N .

Proof:

As every x-c.p.I of N is c.p.I of N (by proposition (2.8)) and every c.p.I of N is c.s.p.I of N (by proposition (1.14)) then every x-c.p.I of N is c.s.p.I of N .

2.10 Proposition:

Let N be a Boolean α_1, α_2 near-ring, then every x-c.s.p.I of N is c.s.p.I of N .

Proof:

As every x-c.s.p.I of N is c.p.I of N (by proposition (2.7)) and every c.p.I of N is c.s.p.I of N (by proposition (1.14)) then every x-c.s.p.I of N is c.s.p.I of N .

2.11 Proposition:


Let N be a Boolean α_1, α_2 near-ring, then every x-c.s.p.I of N is x-c.p.I of N .

Proof:

As every x-c.s.p.I of N is c.p.I of N (by proposition (2.7)) and every c.p.I of N is x-c.p.I of N (by proposition (2.8)) then every x-c.s.p.I of N is x-c.p.I of N .

2.12 Remark:

The following diagram show us the relationships between these ideals



3 Completely and semi completely prime ideal α_1, α_2 near-ring with respect to an element

3.1 definition:

Let N be a α_1, α_2 near-ring, is called a completely semi prime ideal near ring, denoted by (c.s.p.I α_1, α_2 near-ring) if every ideal of N are c.s.p.I of N .

3.2 example:

Let N be a α_1, α_2 near-ring, in example(2.2) is a c.s.p.I of N since all its ideals $I_1=\{0,a\}$, $I_2=\{0,b\}$, $I_3=\{0,a,b\}$, $I_4=\{0\}$, $I_5=N$ are c.s.p.I of N .

3.3 definition:

Let N be a α_1, α_2 near-ring, is called a completely prime ideal near ring, denoted by (c.p.I α_1, α_2 near-ring) if every ideal of N are c.p.I of N .

3.4 example:

Let N be a α_1, α_2 near-ring, in example(2.2) is a c.p.I of N since all its ideals $I_1=\{0,a\}$, $I_2=\{0,b\}$, $I_3=\{0,a,b\}$, $I_4=\{0\}$, $I_5=N$ are c.p.I of N .

3.5 proposition:

Let N be a α_1, α_2 near-ring, if N is a c.p.I α_1, α_2 near-ring, then N is a c.s.p.I α_1, α_2 near-ring.

Proof:

let N is a c.p.I α_1, α_2 near-ring so, every ideal of N is a c.p.I of N then, every ideal of N is a c.s.p.I of N (by proposition(1.14)) then, N is a c.s.p.I α_1, α_2 near-ring.

3.6 definition:

Let N be a α_1, α_2 near-ring, is called a x- completely semi prime ideal α_1, α_2 near-ring, denoted by (x-c.s.p.I α_1, α_2 near-ring) if every ideal of N are x-c.s.p.I of N , where $x \in N$.

3.7 example:

Let N be a α_1, α_2 near-ring, in example(2.2) is a (x-c.s.p.I of N) since all its ideals $I_1=\{0,a\}$, $I_2=\{0,b\}$, $I_3=\{0,a,b\}$, $I_4=\{0\}$, $I_5=N$ are x-c.s.p.I of N .

3.8 definition:

Let N be a α_1, α_2 near-ring is called a x- completely prime ideal α_1, α_2 near-ring, denoted by (x-c.p.I α_1, α_2 near-ring) if every ideal of N are x-c.p.I of N , where $x \in N$.

3.9 example:

Let N be a α_1, α_2 near-ring in example(2.2) is a (x-c.p.I of N) since all its ideals $I_1=\{0,a\}$, $I_2=\{0,b\}$, $I_3=\{0,a,b\}$, $I_4=\{0\}$, $I_5=N$ are x-c.p.I of N .

3.10 proposition:

If N is a x -c.p.l α_1, α_2 near-ring, then N is a x -c.s.p.l α_1, α_2 near-ring, where $x \in N$.

Proof:

Let N is a x -c.p.l α_1, α_2 near-ring so, every ideal of N is a x -c.p.l of α_1, α_2 near-ring then, every ideal of N is a x -c.s.p.l of α_1, α_2 near-ring (by proposition(2.5)) then, N is a x -c.s.p.l α_1, α_2 near-ring.

3.11 proposition:

Let N be a Boolean α_1, α_2 near-ring then N be c.p.l iff N is an x -c.s.p.l for all $x \in N$.

Proof:

\Rightarrow let N is c.p.l of α_1, α_2 near-ring (so, every ideal of N is a c.p.l of N then, every ideal of N is a x -c.s.p.l of α_1, α_2 near-ring (by proposition(2.7)) then, N is an x -c.s.p.l for all $x \in N$ (\Leftarrow) let N is an x -c.s.p.l for all $x \in N$ so, every ideal of N is a x -c.s.p.l of N then, every ideal of N is a c.p.l of N (by proposition(2.7)) then N be c.p.l

3.12 proposition:

Let N be a Boolean α_1, α_2 near-ring then N be c.p.l of N iff N is an x -c.p.l for all $x \in N$.

proof:

Obvious by definitions above and proposition (2.8)

3.13 Proposition:

Let N be a Boolean α_1, α_2 near-ring, if N is x -c.p.l then N is c.s.p.l

Proof:

Obvious by definitions above and proposition (2.9)

3.14 Proposition:

Let N be a Boolean α_1, α_2 near-ring, if N is x -c.s.p.l then N is c.s.p.l

Proof:

Obvious by definitions above and proposition (2.10)

3.15 Proposition:


Let N be a Boolean α_1, α_2 near-ring, if N is x -c.s.p.l then N is x -c.p.l.

Proof:

Obvious by definitions above and proposition (2.11)

3.16 Remark:

The following diagram show us the relationships between these α_1, α_2 near-ring



4 Completely and semi completely prime ideal with respect to an multiplicative identity of a α_1, α_2 near-ring

4.1 proposition:

Let N be a Boolean α_1, α_2 near-ring with multiplicative identity e' then I is e' -c.s.p.l of N iff it is a c.s.p.l of N .

4.2 proposition:

Let N be a Boolean α_1, α_2 near-ring with multiplicative identity e' then I is e' -c.p.l of N iff it is a c.p.l of N .

4.3 proposition:

Let N be a Boolean α_1, α_2 near-ring with multiplicative identity e' then I is e' -c.p.l of N iff it is a e' -c.s.p.l of N .

Proof:

(\Rightarrow) let $\forall y \in N, x.y^2 \in I$
 as I is e' -c.p.l of N
 let $x.y.y \in I$, then
 $e'.x.y.y \in I$, then
 $e'.x.y.y = e'.x.y^2 = x.y^2 \in I$
 then, $y \in I$ as I is e' -c.p.l of N
 so, I is a e' -c.s.p.l of N
 (\Leftarrow) let $\forall y \in N, x.y.y \in I$
 as, I is a e' -c.s.p.l of N
 Let $x.y^2 \in I$, then
 $e'.x.y^2 \in I$, then
 $e'.x.y^2 = e'.x.y.y = x.y.y \in I$
 then, $y \in I$ as I is a e' -c.s.p.l of N
 so, I is e' -c.p.l of N .

4.4 example:

Let $N = \{0, a, b, c\}$ be a α_1, α_2 near-ring, with addition and multiplication defined as:

+	0	a	b	c	.	0	a	b	c
0	0	a	b	c	0	0	0	0	0
a	a	0	c	b	a	0	a	a	0
b	b	c	0	a	b	0	a	b	c
c	c	b	a	0	c	0	0	c	c


the ideal $I = \{0, a\}$ is b -c.p.l of N iff it is a b -c.s.p.l of N , but its same example which clear that not all c.p.l of N are x -c.p.l for all $x \in N$ since $I = \{0, a\}$ is c.p.l of N but it is not a c -c.p.l of N , since $a.(b.c) = 0 \in I$ but $b \notin I$ and $c \notin I$ and not all c.s.p.l of N are x -c.s.p.l of N since $I = \{0, a\}$ is c.s.p.l of N but it is not a c -c.s.p.l of N , since $a.b^2 \in I$ but $b \notin I$.

4.5 proposition:

Let N be a Boolean α_1, α_2 near-ring with multiplicative identity e' then I is c.p.l of N iff it is a e' -c.s.p.l of N .

4.6 Remark:

The following diagram show us the relationships between these ideals



REFERENCES

- [1] G.Birkenmeier,H.Heatherly and E.Lee,"Prime Ideals in Near-Ring",Results Math24(1993),27-48.
- [2] G .Pilz"Near-Ring",North-Holland Publishing Company, Amsterdam-New York –oxford, 1983.
- [3] Husein Hadi Abbass, Mohanad Ali Mohammed "On completely Prime Ideal with respect to an element of a Near-Ring",Journal of kerbala university , vol. 10 no.3 scientific . 2012,285-301 .
- [4] Hussien Hadi Abass, Showq Mhammed .E"Completely Semi Prime Ideal With Respect to an Element of a Near Ring",Journal of Kufa for Mathematics and Computer, Vol.1, No.3, may , 2011, 1- 7 .
- [5] H.A.S.Abujabal,"On structure and commutativity of Near-Rings",Antofagasta-Chile,2000.
- [6] N.Groenewald,"The completely prime radical in Near-Rings",Acta Math.Hung.51(1988),301-305.
- [7] N.J.Groenewald,"Different prime ideals in Near-Ring",comm.Algebra,19(10)(1991),2667-2675.
- [8] S.uma,R.Balakrishnam,T.Tamizh" α_1, α_2 near-ring international journal of algebra,vol.4,2010 ,no.2,71-79.
- [9] W. B. Vasantha Kandasamy"smarandache Near-Rings",America Research Press and W.B. Vasantha Kandasamy,2002,19-23.
- [10] W.M.L.Holcombe,"Primitive Near-Ring",doctoral Dissertation,university of Leeds,1970.